

Nutritional Values and Therapeutic Uses of *Capra hircus* Milk

Gaurav Tiwari¹, Akanksha Chauhan¹, Pooja Sharma², Ruchi Tiwari^{1*}

¹PSIT-Pranveer Singh Institute of Technology (Pharmacy), Kanpur - Agra - Delhi, NH2, Bhauti, Kanpur, Uttar Pradesh, INDIA.

²Department of Pharmacy, Dr. Bhimrao Ambedkar University, Paliwal Park, Park Rd, Agra, Uttar Pradesh, INDIA.

ABSTRACT

Capra hircus (Goats) are a vital part of the livestock industry because of their ability to adapt to harsh conditions, making them ideal for landless and marginal farmers. The *Capra hircus* is one of the most prominent options of milk and meat for humans. In terms of structure and wholesome qualities, *Capra hircus* milk differs from those of dairy animals and humans. *Capra hircus* make a major contribution to the supply of milk and milk products, as well as to the rural economy and wellbeing. *Capra hircus* are positioned seventh on the planet regarding milk creation from various species. The calcium, magnesium, and phosphorus substance in *Capra hircus* milk is more prominent than that of cow or human drain. *Capra hircus* milk proteins and medium-chain fatty substances have been portrayed as particular lipids and proteins with unmistakable medical advantages. A few bioactive compounds display in *Capra hircus* milk and other *Capra hircus* -derived products have been shown to benefit patients suffering from several chronic illnesses. Goat's milk contains many peptides, fats, and oligosaccharides that can help with cardiovascular disease, metabolic

disorders, neurological degeneration, and intestinal health. They've also been found to have cancer-preventive properties. Furthermore, the oligosaccharides used in goat's milk have immunomodulatory properties, inhibit pathogenic bacteria adhesion, and have prebiotic and pro-bifidogenic properties. *Capra hircus* milk is particularly recommended for children, older adults, and people recovering from illness due to its possible health benefits.

Keywords: *Capra hircus* milk's compositions, *Capra hircus* milk's products, Medicinal values of *Capra hircus* milk.

Correspondence

Prof. Ruchi Tiwari

PSIT-Pranveer Singh Institute of Technology (Pharmacy), Kanpur - Agra - Delhi, NH2, Bhauti, Kanpur-209305, Uttar Pradesh, INDIA.

Email id: tiwaridrruchi@gmail.com

DOI: 10.5530/ijpi.2022.4.71

INTRODUCTION

The milk contains few physiologically useful segments including the proteins, nutrients, for an example, the vitamin E, and the vitamin C just as carotenoids and flavonoids with the cell reinforcement properties. Thusly, milk with a higher cell reinforcement limit will reflect the more noteworthy oxidative security and give conceivably more prominent insurance to the purchaser from openness to the oxidative pressure that is perceived as a noticeable component of numerous intense and persistent infections.^{1,2} Lately, the significant interest has focused on goat's milk for its nourishing and biofunctional properties.³ *Capra hircus* were among the primary livestock to be domesticated. As shown by the archaeological evidence, they have been related to the man in a harmonious relationship for as long as 10,000 years.⁴ In many parts of the world, *Capra hircus* are significant milk producers of the tropics and make a significant contribution a significant contribution to human nutrition in the majority of developed countries.⁵ A *Capra hircus* is often referred to as the "poor man's cow."⁶ *Capra hircus* are available altogether of the landmasses and the world's absolute quantities of *Capra hircus* were 861.9 million. The biggest number of *Capra hircus* is seen in Asia, trailed by Africa, in Asia 514.4 million, Africa 291.1 million, Northern America 3.0 million, Central America 9.0 million, Caribbean 3.9 million, South America 21.4 million, Europe 18.0 million and Oceania 0.9 million.⁷ Around 80% of *Capra hircus* milk is produced in Asia.⁶ In such a manner primary nation incorporate India, Bangladesh, Pakistan, China, Iran and Turkey. These Figures demonstrate are a solid sign of the significance of *Capra hircus* in the livelihoods of individuals in creating and immature countries, especially in Asia.⁸ The *Capra hircus* sector has inflated considerably throughout the last decades. From 1980 until 1999 the quantities of *Capra hircus* farms have ranch have expanded by 55%

and the *Capra hircus* milk creation by 58%. In any case, these numbers are probably going to be a lot more significant, given because of the lot of unreported home utilization particularly in developed countries.⁹ Cow milk creation is less expensive and the volumes are much larger than cow milk has a lower market value. Commercial *Capra hircus* milk creation is more costly due to bring down profitability, differences due to the season, and therefore they would like of larger animal groups. *Capra hircus* milk varies from cow milk in terms of digestibility, alkalinity, buffering limit, and certain therapeutic values of medicine and human nutrition, in addition to manufacturing variations. Goat's milk and determined dairy items expect significance in the human eating regimen with specific pertinence in babies for whom goat's milk addresses an option in a contrast to cow's milk.¹⁰ From a healthful perspective, goat's milk is unique to concerning that of different species. It is characterized by little measured unsaturated fat globules and a high substance of short-chain unsaturated fats which improve fat edibility and internal organ absorption compared with cow's milk. The fat globules in *Capra hircus* milk were bountiful in sizes under 3.5 µm contrasted with a cow which was 4.55 µm. A few investigations revealed that the quantity of fat globules less than 5 µm in *Capra hircus* milk is ~80% contrasted with ~60% in cow milk. This trademark adds to the gentler surface of *Capra hircus* milk items and upgrades lipid digestion, along these lines making them more absorbable.¹¹ Also, the presence of medium-chain unsaturated fats (caproic, caprilic, and capric corrosive) in *Capra hircus*' milk has been credited to a decrease of cholesterol in human tissues by restricting cholesterol stockpiling and improving its preparation.¹² Based on the properties of αS1-casein, Goat's milk is believed to have a lower allergenic capacity than cow's milk.¹³ The fat in *Capra hircus* milk

Copyright © 2022 Author(s). Exclusive Licensee Phcog.Net. Distributed under a Creative Commons Attribution License (CC BY 4.0).

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

contains a higher extent of medium-chain unsaturated fats and formed linoleic acid, which is related to the qualities of cheddar flavor and “goaty” smell of *Capra hircus* milk just as the counter anti-carcinogenic and anti-atherogenic effect. Other than fat, protein in *Capra hircus* milk has a larger buffering limit and particular alkalinity, which might be helpful for the treatment of stomach ulcers. This advantage is because of more elevated levels of major buffering segments, like proteins, non-protein nitrogen, and phosphate.¹¹ Another distinctive characteristic of *Capra hircus* milk is casein micelles. They are less solvated, have lower heat stability, and lose -casein more quickly than cow milk. They were also crucial in the cheese-making process, especially during the edge.¹² In a new report it has been shown that every day utilization of *Capra hircus* milk in the eating regimen of immuno-bargained matured patients goes about as a down regulator of intense irritation, decreasing the misrepresented basal discharge of interleukin (IL)- 8 and IL-6 intense reaction, and applying a moderate down-guideline of IL-1b and tumor necrosis factor (TNF)- a creation. Antithrombotic properties of casein macro-peptides from bovine, ovine, and caprine milk are demonstrated by the movement of the g-fibrinogen 400e411 peptide.¹³ Milk content is determined by milk composition, which varies depending on breed, age, body size, weight, udder size, diet, lactation stage, season, dry period length, and environmental temperature.¹⁴ *Capra hircus* items (mainly meat and dairy) have complex taste, aromas, and leanness, as well as a specific composition of fats, proteins, amino acids, and fatty acids, and have been served in many other nations of the world for decades.¹⁵ Moreover, the beneficial properties of *Capra hircus* milk, as wells as its lower allergenicity when compared to cow milk, has started the interest in *Capra hircus* milk as a nutritional food and it is presently one of the latest things in good dieting in created nations.¹⁶ Also, the utilization of the milk with specific nutritional properties, alone or in a mix with bacterial strains that also have probiotic properties and also can contain physiologically active metabolites, maybe one of the options for producing new dairy functional beverages.¹⁷ Many minerals, like iron, calcium, phosphorous, and magnesium, are better used by *Capra hircus* milk and milk products in the digestive and metabolic processes.¹⁸ Since platelet transfusion is not always possible from outside, *Capra hircus* milk is commonly prescribed to dengue patients to maintain body fluid balance. *Capra hircus* milk diet increased biliary cholesterol secretion, lowering plasma cholesterol levels, but phospholipids, biliary acid, and lithogenic levels remained unchanged.¹⁹

Difference between *Capra hircus* Milk vs. Cow Milk

In several respects, *Capra hircus* milk is a very healthy food, because it is naturally homogenized. *Capra hircus* milk will be much easier to digest than cow milk, as it contains fewer protein molecules than cow milk. *Capra hircus* milk fat molecules have a smaller, more fragile membrane, about half the size of cow milk fat molecules. In the Table 1, the differences in constituents between cow and *Capra hircus* milk are mentioned.²⁰

Beneficial Health Effects of Goat's Milk over the Cow's Milk

When *Capra hircus* milk is compared to cow milk, it has been discovered that *Capra hircus* milk aids in the digestion and metabolic utilization of many minerals such as iron, phosphorous, calcium, and magnesium, as well as the prevention of diseases such as anemia and bone demineralization. There was a greater recovery with *Capra hircus* milk in cases of ferrocene anemia and bone demineralization. *Capra hircus* milk has a beneficial effect on mineral metabolism. Patients who are allergic to cow milk may also use it as a supplement.²¹ *Capra hircus* milk contains a high concentration of fat globules that are smaller than those found in cow milk; globule diameters average about 3.6 and 3.0 m, respectively, compared to 4.0 m in cow milk (Table 2). *Capra hircus*-derived goods have a finer texture due to the smaller size of fat globules.

Table 1: Constituents difference between cow and *Capra hircus* milk.^[24]

Constituents	Cow Milk	<i>Capra hircus</i> Milk
Protein (%)	3.2	3.4
Lactose (%)	4.7	4.1
Vitamin B1 (mg)	–	0.05
Vitamin A (I.U)	158	120
Ash (%)	0.71	0.77
Fat (%)	3.6	3.8

Table 2: Table showing composition of *Capra hircus* milk.^[27]

Compositions	<i>Capra hircus</i> Milk
Energy (kcal)	70
Protein (%)	3.2
Fat (%)	4.0-4.5
Water (%)	87.5
Lactose (%)	4.6
Total Solids (g)	12.2
Ash (g)	0.8

Capra hircus milk also has a lower concentration of 1-casein, which gives it a lower viscosity and higher water-holding capacity.²² Because of these benefits, the taste of Goat's milk is distinct and more intense than that of cow's milk, which can limit consumer acceptance of its derivatives. *Capra hircus* milk, on the other hand, is easier to absorb than cow milk because it lacks agglutinins (Table 3).²³

Composition/ Nutritional Values of *Capra hircus* milk

Dairy food products derived from milk are considered to be an integral part of a well-balanced diet. Milk, like a mammal's first meal, provides all of the energy and supplements required for the neonate's proper growth and development. Milk intake ceases in all mammals after weaning, except for humans, who continue to drink milk throughout their lives. It contains more fat, protein, and ash than cow milk, but less lactose. *Capra hircus* milk has a lower overall casein content but a greater nonprotein nitrogen content than cow milk. *Capra hircus* and cow milk has 3 to 4 times the protein and ash content of human milk. *Capra hircus*, cow, and human milk have identical total solids and caloric values.²⁴ Milk has different physical characteristics and structures depending on the species. Milk is a complicated oil-in-water emulsion that includes proteins, fats, carbohydrates (primarily lactose), and trace minerals, enzymes, immunoglobulins, hormones, cells, and vitamins.¹²

Lipids

In terms of expense, nutrition, physical, and sensory characteristics that dairy products impart, lipids are the most important components of *Capra hircus* milk.²⁵ Triacylglycerol (TAG), which contains a higher number of esterified fatty acids, makes up about 97% of the lipid fraction of *Capra hircus* milk.²⁷ Easy lipids like diacylglycerols, monoacylglycerols, and cholesterol esters, as well as complex lipids like phospholipids and liposoluble compounds like sterols, cholesterol esters, and hydrocarbons, are both included in the lipid fraction.²⁷ *Capra hircus* milk fat content varies between breeds, ranging from 2.45 to 7.76 percent. Fat globule diameters in *Capra hircus*, sheep, buffalo, and cow milk were found to be 3.49, 4.55, 5.92, and 3.30 m, respectively.²⁸ Tinier fat globules allow for better fat dispersion and homogeneity in *Capra hircus* milk, resulting in a larger surface area of fat for improved digestion by lipases.²⁹ *Capra*

Table 3: Table showing derivatives of *Capra hircus* and its applications.

Derivatives of <i>Capra hircus</i> Milk	Therapeutic effects of <i>Capra hircus</i> Milk	Use of <i>Capra hircus</i> Milk	References
Bioactive peptides	Hypertension	With individual proteases such as thermolysin, trypsin, subtilisin, papain, and pepsin or their combinations, the bioactivity of <i>Capra hircus</i> milk protein hydrolysates and the release of ACE-inhibitory and anti-oxidant peptides	[27,28]
Antimicrobial Peptides	Anti-inflammatory, Anti-osteoporotic	Alpha-S2 casein is present in Ethawah <i>Capra hircus</i> milk and yogurt. This protein contains eight bioactive peptides with various effects, including anti-osteoporotic and anti-inflammatory properties, and it was not present in cow fresh milk.	[24,30]
Cytomodulatory and anticancer peptides	Anti-cancer	Lactoferrin from <i>Capra hircus</i> milk not only has antimicrobial properties, but it also induces apoptosis in a human cervical cancer cell line. Given its similarity to other lactoferrins, <i>Capra hircus</i> lactoferrin may be able to provide protection against other cancers.	[28,32]
Antioxidant peptides	Prevent or delay oxidative stress	milk proteins just as milk-inferred proteins have been considered as likely transporters for the conveyance of antioxidant prevention agent peptides in the gastrointestinal plot, where they may apply direct defensive impacts by searching receptive oxygen species and decreasing the oxidative pressure	[23,31]
Immunomodulatory/ anti-inflammatory peptides	Anti-inflammatory	Colostrinin (CLN) is a proline-rich polypeptide complex isolated from ovine colostrum. Anti-inflammatory and antioxidant properties of colostrinin Colostrinin also prevents inflammatory conditions in neurons caused by -amyloid accumulation.	[24,25]
Prevention of milk allergy	Allergy	Cow's milk allergy is a common pediatric condition with an incidence of around 2.5 % within the first three years of life. It's an IgE-mediated allergy, which means the body develops antibodies to certain proteins (allergens) in cow milk.	[23,31]
Prebiotic and antipathogenic activity		The optimal development of the immune system relies on intestinal colonization with a healthy microbiota, and there is a lot of research and commercial interest in adjusting the microbiota for health purposes.	[28,34]
Anti-inflammatory activity	Anti-inflammation	In the newborn child, <i>Capra hircus's</i> milk oligosaccharides can serve as anti-inflammatory agents.	[23,37]
Prevention of inflammatory bowel disease (IBD) and colitis	Bowel disease	The healing mechanism after DSS-induced colitis is helped by <i>Capra hircus</i> milk oligosaccharides. Furthermore, oligosaccharides containing N-acetylglucosamine, which promotes <i>B. bifidum</i> formation, can stimulate the development of the intestinal flora.	[28,39]
Bioactive peptides	Hypertension	With individual proteases such as thermolysin, trypsin, subtilisin, papain, and pepsin or their combinations, the bioactivity of <i>Capra hircus</i> milk protein hydrolysates and the release of ACE-inhibitory and anti-oxidant peptides	[28,24]
Antimicrobial Peptides	Anti-inflammatory, Anti-osteoporotic	Alpha-S2 casein is present in Ethawah <i>Capra hircus</i> milk and yogurt. This protein contains eight bioactive peptides with various effects, including anti-osteoporotic and anti-inflammatory properties, and it was not present in cow fresh milk.	[24]
Cytomodulatory and anticancer peptides	Anti-cancer	Lactoferrin from <i>Capra hircus</i> milk not only has antimicrobial properties, but it also induces apoptosis in a human cervical cancer cell line. Given its similarity to other lactoferrins, <i>Capra hircus</i> lactoferrin may be able to provide protection against other cancers.	[28]
Antioxidant peptides	Prevent or delay oxidative stress	milk proteins just as milk-inferred proteins have been considered as likely transporters for the conveyance of antioxidant prevention agent peptides in the gastrointestinal plot, where they may apply direct defensive impacts by searching receptive oxygen species and decreasing the oxidative pressure	[23]
Immunomodulatory/ anti-inflammatory peptides	Anti-inflammatory	Colostrinin (CLN) is a proline-rich polypeptide complex isolated from ovine colostrum. Anti-inflammatory and antioxidant properties of colostrinin Colostrinin also prevents inflammatory conditions in neurons caused by -amyloid accumulation.	[24]
Prevention of milk allergy	Allergy	Cow's milk allergy is a common pediatric condition with an incidence of around 2.5 % within the first three years of life. It's an IgE-mediated allergy, which means the body develops antibodies to certain proteins (allergens) in cow milk.	[20]
Prebiotic and antipathogenic activity		The optimal development of the immune system relies on intestinal colonization with a healthy microbiota, and there is a lot of research and commercial interest in adjusting the microbiota for health purposes.	[28,34]
Anti-inflammatory activity	Anti-inflammation	In the newborn child, <i>Capra hircus's</i> milk oligosaccharides can serve as anti-inflammatory agents.	[23,35]
Prevention of inflammatory bowel disease (IBD) and colitis	Bowel disease	The healing mechanism after DSS-induced colitis is helped by <i>Capra hircus</i> milk oligosaccharides. Furthermore, oligosaccharides containing N-acetylglucosamine, which promotes <i>B. bifidum</i> formation, can stimulate the development of the intestinal flora.	[28,32]

continued...

Table 3: Cont'd.

Derivatives of <i>Capra hircus</i> Milk	Therapeutic effects of <i>Capra hircus</i> Milk	Use of <i>Capra hircus</i> Milk	References
Bioactive peptides	Hypertension	With individual proteases such as thermolysin, trypsin, subtilisin, papain, and pepsin or their combinations, the bioactivity of <i>Capra hircus</i> milk protein hydrolysates and the release of ACE-inhibitory and anti-oxidant peptides	[28]
Antimicrobial Peptides	Anti-inflammatory, Anti-osteoporotic	Alpha-S2 casein is present in Ethawah <i>Capra hircus</i> milk and yogurt. This protein contains eight bioactive peptides with various effects, including anti-osteoporotic and anti-inflammatory properties, and it was not present in cow fresh milk.	[24]
Cytomodulatory and anticancer peptides	Anti-cancer	Lactoferrin from <i>Capra hircus</i> milk not only has antimicrobial properties, but it also induces apoptosis in a human cervical cancer cell line. Given its similarity to other lactoferrins, <i>Capra hircus</i> lactoferrin may be able to provide protection against other cancers.	[28]
Antioxidant peptides	Prevent or delay oxidative stress	milk proteins just as milk-inferred proteins have been considered as likely transporters for the conveyance of antioxidant prevention agent peptides in the gastrointestinal plot, where they may apply direct defensive impacts by searching receptive oxygen species and decreasing the oxidative pressure	[23]
Immunomodulatory/ anti-inflammatory peptides	Anti-inflammatory	Colostrin (CLN) is a proline-rich polypeptide complex isolated from ovine colostrum. Anti-inflammatory and antioxidant properties of colostrin Colostrin also prevents inflammatory conditions in neurons caused by -amyloid accumulation.	[24,41]
Prevention of milk allergy	Allergy	Cow's milk allergy is a common pediatric condition with an incidence of around 2.5 % within the first three years of life. It's an IgE-mediated allergy, which means the body develops antibodies to certain proteins (allergens) in cow milk.	[42]
Prebiotic and antipathogenic activity		The optimal development of the immune system relies on intestinal colonization with a healthy microbiota, and there is a lot of research and commercial interest in adjusting the microbiota for health purposes.	[28]
Anti-inflammatory activity	Anti-inflammation	In the newborn child, <i>Capra hircus's</i> milk oligosaccharides can serve as anti-inflammatory agents.	[23,39]
Prevention of inflammatory bowel disease (IBD) and colitis	Bowel disease	The healing mechanism after DSS-induced colitis is helped by <i>Capra hircus</i> milk oligosaccharides. Furthermore, oligosaccharides containing N-acetylglucosamine, which promotes <i>B. bifidum</i> formation, can stimulate the development of the intestinal flora.	[28]

hircus milk fat contains 97-99 percent free lipids (roughly 97 percent of which are triglycerides) and 1-3 percent bound lipids (about 47 percent neutral and 53 percent polar lipids). Short- and medium-chain-length fatty acids (MCT) are substantially higher in *Capra hircus* milk fat than in cow and human milk. This property has been used to help patients with several fat malabsorption issues.^{23,25,26,30} *Capra hircus* milk also contains more polyunsaturated fatty acids (PUFA) and conjugated linoleic acid (CLA). Short and medium-chain fatty acids, as well as medium-chain triacylglycerols (MCT), have become well-known prescription therapies for a variety of clinical conditions. The MCTs are consumed in their entirety in the intestine and are not degraded or re-esterified. Since the molecules are taken up directly into the portal vein, no micelle formation is needed for absorption.^{21,31} Minor lipids present in *Capra hircus* milk include glycolipids, gangliosides, glycosphingolipids, and cerebroside. These minor lipids, which are also bioactive components of *Capra hircus* milk, have effects that are close to those in cow and human milk. Cell-to-cell communication, immune recognition, and receptor roles for protein hormones and bacterial toxins including enterotoxin and cholera toxin are among these functions.³¹

Proteins

Milk proteins are divided into two phases: an unstable micellar phase made up of casein and a soluble phase made up of whey proteins. The main whey proteins are α -lactoglobulin and α -lactalbumin, and the antigenicity of β -lactoglobulin can be partly eliminated by some therapies. Caseins make up about 80% of the proteins and are known as α s1, α s2, β and k-caseins.²⁸ As compared to cow milk, *Capra hircus* milk has lower levels of α s-casein, higher amounts of β -casein fractions,

and almost equivalent amounts of k-casein fractions. α S1-casein is the main protein in cow milk, while β -casein is the main protein in *Capra hircus* milk. α S1-casein is found in *Capra hircus* milk, but the amount and genetic variations vary depending on the *Capra hircus* population.²⁹ Differences in digestibility are caused by various genetic groups. Since α 1-casein is only partially digested by gastric juice enzymes, the more α s1-casein in milk, the

longer digestion takes. It is not completely hydrolyzed until it comes into contact with the enzymes in the duodenum.³¹ In *Capra hircus* milk, the concentrations of folate-binding protein are higher than in cow milk. Since the protein has a strong affinity for binding folate, it prevents humans from digesting and absorbing it. *Capra hircus* milk is a total protein source, containing all essential amino acids without the fat and mucus-producing components found in cow milk.²⁷ Immunoglobulin, lactoperoxidase, lactoferrin, folate binding protein, and, more recently, α -lactalbumin and β -lactoglobulin are examples of whey and casein milk proteins, as well as their bioactive peptides. The nutritional benefits of milk proteins are determined by the number of essential amino acids present. Milk amino acid levels per 100 g of protein vary only slightly between species, which is most likely due to variations in total protein content. *Capra hircus* milk contains more essential amino acids than cow milk, including threonine, lysine, leucine, tyrosine, cysteine, phenylalanine, valine, and nonessential proline and glutamic acid.³¹

Carbohydrates

Capra hircus milk's main carbohydrate is lactose, which is significantly lower in quality than cow milk. It is made in the mammary gland from glucose and galactose, with the milk protein α -lactalbumin

playing a key role. Lactose is an essential nutrient since it helps in the absorption of calcium, magnesium, and phosphorus in the intestine, as well as the utilization of vitamin D. It's also crucial for milk synthesis and milk secretion into the udder's duct system [11]. Oligosaccharides, glycopeptides, glycoproteins, and nucleotides are small quantities of carbohydrates contained in *Capra hircus* milk. When compared to cow milk, *Capra hircus* milk has a higher concentration of lactose-derived oligosaccharides.²⁹ Because of their prebiotic and anti-infective properties, milk oligosaccharides are thought to be beneficial to human nutrition.²¹ *Capra hircus* milk oligosaccharides have been shown to have anti-inflammatory properties in induced colitis in animal models. These findings can help in the treatment of inflammatory bowel disease. Nucleotides in milk are also interesting because they are glycosyl donors for glycosyl transferase in milk and the mammary gland, as well as precursors of glycoproteins, glycolipids, and oligosaccharides in milk biosynthesis. *Capra hircus* milk contains a surprising number of nucleotides. In contrast to human milk, *Capra hircus* milk has a higher concentration of oligosaccharides than bovine and ovine milk, and the oligosaccharide structures contained in *Capra hircus* milk are the most similar to those found in human milk. This is especially important for infant feeding, as human milk oligosaccharides have prebiotic and anti-infective properties that are extremely beneficial to infants. Increased butyrate synthesis and the reduction of pro-inflammatory bacterial species by inhibiting their adhesion to the epithelial membrane, reducing bacterial translocation, and promoting selective growth of beneficial *Lactobacillus* and *Bifidobacteria* species are the mechanisms behind their anti-inflammatory impact.²⁴

Vitamins

Since *Capra hircus* turn all β -carotene from foods into vitamin A in the milk, *Capra hircus* milk has a higher vitamin A content than cow milk. *Capra hircus* milk is often whiter than cow milk for the same reason. Vitamin B6 and Vitamin D, which are both essential during infancy, are low in both *Capra hircus* and cow milk. Vitamin A levels in *Capra hircus* milk are similar to those in human milk. Vitamin A is necessary for innate and adaptive immune responses, such as cell-mediated immunity and antibody responses. This vitamin deficiency has been linked to both increased and decreased innate immunity, affecting NK cell function and phagocytic activity. In addition to its function in bone health maintenance, vitamin D plays an important role in the immune system and may help prevent infections, autoimmune diseases, cancer, and diabetes. Vitamin D deficiency appears to impair NK cell activity, resulting in a reduction in infections. Eventually, vitamin C is a well-known water-soluble antioxidant that is widely available in *Capra hircus* milk compared to cow milk. Antiviral and anti-oxidant properties of this vitamin have been shown to affect many facets of the immune system, including immunity control.²⁵

Minerals

Potassium, Calcium, Chloride, Phosphorus, Selenium, Zinc, and Copper are all said to be higher in *Capra hircus* milk than in cow milk. It's important to note, however, that *Capra hircus* feed can have a significant impact on the mineral content of the milk. Potassium is essential for acid/base balance, as well as muscle, nerve, and kidney function. Fluid equilibrium, blood H, and osmotic pressure are all maintained by chloride. It also has an impact on liver function and is a major component of hydrochloric acid in the stomach's gastric juice. Calcium is essential for bone formation, but it also affects muscle function, nerve function, and blood coagulation. Phosphorous, like calcium, is essential for bone structure, muscle function, and nerve function, but it also plays a role in energy production. Selenium aids in the defense of cells against free radicals. It has a beneficial effect on certain immune

cells (leukocytes) and protects the body from heavy metals. Zinc is an essential component of many enzymes involved in carbon dioxide transport, protein synthesis, and carbohydrate metabolism. It also works with the hormone insulin to control carbohydrate metabolism. Many enzymes need copper as well. It affects iron and oxygen metabolism, as well as cell defense against free radicals. Iodine is required for two thyroid enzymes that regulate metabolism and promote body growth and development. In comparison to human milk, *Capra hircus* milk has a higher zinc content. Zinc is a mineral that is essential for maintaining healthy skin, wound healing, and is involved in both innate and adaptive immunity. Zinc also has antioxidant properties and acts as a cofactor for the antioxidant enzyme superoxide dismutase, which helps to remove reactive oxygen species. Similarly, selenium is an important mineral for immune health since it affects both innate and adaptive immunity. Selenium is a necessary cofactor for the antioxidant enzyme glutathione peroxidase, which is responsible for scavenging harmful free radicals in the body and activating macrophages.^{23,27}

Capra hircus-derived products and their nutritional values

The use of *Capra hircus* items was initially connected to a range of medical issues, including food allergies to cow milk proteins. During the first three years of life, cow milk allergy is fairly normal. According to several reports, *Capra hircus* milk can help with between 30 and 40% of cases.¹¹ *Capra hircus* and cows have a variety of physiological and anatomical differences that influence the composition of *Capra hircus* milk and its products (Figure 1).¹⁰

Capra hircus milk items are known to be the most marketable dairy products. As a result, many aspects of *Capra hircus* milk are currently the subject of increased analysis.²⁷ Two glasses of *Capra hircus* milk, or the equivalent amount of cheese or yogurt, can provide up to 94% of an adult's recommended dietary allowance (RDA) of essential amino acids, 83% of calcium, and 78 % of riboflavin requirements, while also providing a dietary source of other minerals and vitamins, but to a lesser degree.¹⁰

Capra hircus Yoghurt

Capra hircus yogurt is produced using similar methods to cow milk yogurt, but it has distinct organoleptic properties and nutritional composition. During storage, it has less viscosity, a softer consistency taste, and higher acidity.¹⁷ Free caproic, caprylic, lauric, and myristic




Figure 1: Goat Products and their effect on Immune system.

acids can be found in *Capra hircus* yogurt. Nevertheless, As compared to cow yogurt, palmitic and stearic acids were nearly identical, while oleic, linolenic, and palmitic acids were lower.²⁷

Capra hircus butter/ Ghee

Butter made from *Capra hircus* milk is not widely available, and it is sometimes chemically colored to resemble cow butter. Cream separation is difficult, the texture is smoother, and it has a strong propensity to hydrolytic rancidity.²⁸ *Capra hircus* milk presents a separation problem due to its small fat globules, and it's typical! the processing of ghee made from *Capra hircus* milk is also greasy. As a result, mixing *Capra hircus* milk and buffalo milk in a 1:1 ratio produces high-quality ghee.²⁹

Capra hircus Cheese

Cheese made from *Capra hircus* milk has been made for hundreds of years. In recent years, the production of *Capra hircus* milk cheese has gained a commercial advantage in many Western European countries, since regulation for this form of milk and its products is less restrictive than for cow milk products.³⁰ The characteristics of the milk used in cheese production have a direct influence on its structure and characteristics.³² Milk from *Capra hircus* Gouda cheese is usually made in artisanal units using conventional technology that has been passed down from generation to generation and has a distinct flavor and taste.^{21,23}

Nutrient Functionality

Nutrients have recently become a hot topic in the food and nutrition sciences because they appear to be able to modulate human inflammatory status.²⁴ Milk's ability to support the development of the newborn's immune system, inhibit bacterial growth, and provide anti-oxidative and anti-inflammatory protection makes dairy products a particularly interesting food type to study in the context of inflammation.¹² Different bioactive components of milk and dairy products have been shown to inhibit low-grade systemic inflammation and act as coadjuvants in traditional therapies. Following a brief description of the immune system's structure and function, we will focus on the effects of many bioactive compounds contained primarily in *Capra hircus* milk and their effects on low-grade systemic inflammatory diseases and immunity in the following sections.

Medicinal Properties of Capra hircus Milk (Figure 2)

Antimicrobial Properties

Antimicrobial peptides have also been shown to be precursors to milk proteins. Lactoferrin-derived peptides are the most well-known peptides. Lactoferrin is an iron-binding protein whose primary role is to transport iron. It's involved in several bodily functions, including iron absorption and immune responses. It also acts as an antioxidant and has anti-carcinogenic and anti-inflammatory properties.²⁸ α_2 casein that has been degraded by the gastrointestinal enzyme pepsin has also been shown to have antimicrobial activity.^{23,31} During *Capra hircus* milk fermentation, the pH of the milk decreased more quickly, and a higher number of viable *Bifidobacterium longum* Bb-12 cells were discovered. The higher aging action of lactic corrosive microscopic organisms in *Capra hircus* drain, concurring to a few creators.²⁶

Due to its special composition and structure. Since *bifid bacteria* develop better in the presence of higher levels of certain amino acids found in *lacto globulins* and *lacto albumins*, higher whey protein content may be important.²⁷ *In-vitro* studies have shown that fermented milk containing probiotics inhibits gram-negative bacteria including *Yersinia enterocolitica*, *Escherichia coli*, *Aeromonashydrophila*, and *Salmonella species*.^{28,29} While *Capra hircus* milk has been shown to have strong antibacterial and immunological properties, little is known about the




Figure 2: Effects of Goat milk on different systems of body.

impact of fermented *Capra hircus* milk on pathogenic and potentially pathogenic microorganisms. Fermented *Capra hircus* milk had a greater inhibitory effect on the growth of *Serratia marcescens* colonies than fermented cow milk. *Capra hircus* milk has a particular antimicrobial effect and its particular arrangement may bring about the expanded antimicrobial mixtures creation.^{23,32} Control of pathogenic and spoilage microbes in *Lactobacillus* strains, an essential dairy culture starter variety of foods, is important to ensure food quality and protection when used to make fermented foods.²¹ Biopreservation has as of late been a hot subject. Bacteriocins' disclosure opened up an entire modern world of conceivable outcomes.²² As an alternative to food dyeing, this method is used.^{27,29} Many studies on improving food safety by using natural microflora and bacteriocins developed by lactic acid bacteria isolated from *Capra hircus* milk have been published in recent years in chemical additives for increasing self-life storage. Most studies, however, focus on bacteriocins, which are thought to be healthy since they have long been formed by numerous *lactococci*, *pediococci*, and *leuconostoc*, which are known to be part of the natural flora in fermented foods.^{23,33}

Treatment of Cardiovascular Diseases

In created nations, cardiovascular illness (CVD) is the driving cause of passing. Coronary heart disease, high blood pressure, arrhythmias, and atherosclerosis are only a few of the illnesses that affect the heart and blood vessels. The key cause of CVD is the formation of atherosclerotic plaque in blood vessels, which ultimately leads to a cardiovascular event. The variables that impact the event of a cardiovascular occasion. Tall blood weight, dyslipidemia, diabetes, and corpulence are too chance components.²⁴ Atherosclerosis is a slow-moving disease that may begin in childhood. The exact cause of the disease is unclear, but atherogenic lipoprotein accumulation in the arterial walls appears to be a significant mechanism. Low-density lipoprotein (LDL), also known as "poor cholesterol," is an atherogenic lipoprotein that transports cholesterol from the liver to the blood vessels. The "good" cholesterol is high-density lipoprotein (HDL), which transfers cholesterol from the arteries to the oxidative modification of LDL (ox-LDL), which is essential in the development of atherosclerosis.³⁴ As a result, antioxidants that can prevent LDL oxidation may be useful in preventing atherosclerosis. Medium-chain triglycerides (MCT) found in *Capra hircus* milk include fatty acid esters of caproic, caprylic, and capric fatty acids. In rat models, these MCT have been shown to reduce plasma cholesterol.¹² and to suppress and/or restrict cholesterol deposition in the tissues²⁵ (Figure 3). MCT present in *Capra hircus* milk has anti-atherogenic properties. Cells




Figure 3: Role of Goat milk in blood pressure regulation. ACE: Angiotensin Converting Enzyme, AngI: Angiotensin I, AngII: Angiotensin II.

removed from humans that had been consuming *Capra hircus* milk from various breeds were activated to release nitric oxide by components in the milk, according to an *in-vitro* analysis. Nitric oxide (NO) then travels through the lymphatic system to the bloodstream, where it causes vasodilation and has a cardio-protective and anti-atherogenic effect.^{29,35} Zn bioavailability, an antioxidant mineral, is improved by *Capra hircus* milk. The better nutritive use of *Capra hircus* milk fat,¹² provides a lower substrate for lipid peroxidation and, as a result, reduces the production of free radicals in this form of milk. The favorable impact of *Capra hircus* milk intake on genomic stability, also during Fe-overloading feeding regimes, may be attributed, at least in part, to the high bioavailability of Mg and Zn, as well as its superior fat content.^{12,27}

Instead of the following, magnesium metabolism improves genomic stability: Environmental mutagens and endogenous mechanisms continually disrupt DNA. Cells have developed various forms of DNA repair mechanisms to keep mutation frequencies low. Magnesium is an important cofactor in nearly all stages of nucleotide excision repair and is primarily involved in the removal of DNA damage caused by environmental mutagens. Second, base excision repair is primarily used to repair endogenous DNA damage. The high content of *Capra hircus* milk fat is thought to add to its DNA stabilization benefits. It contains more carnitine, which joins the mitochondria and increases the rate of β -oxidation and energy output from fatty acids, reducing the substrate for lipid peroxidation and, as a result, free radical production.¹⁵

Treatment of Gastrointestinal Diseases

Transmural inflammation characterizes this inflammatory disorder, which may also include ulcerative colitis and Crohn's disease. Inflammatory bowel disease (IBD) is characterized by persistent and relapsing bowel inflammation, but it differs significantly from Crohn's disease in terms of pathophysiology and care. Ulcerative colitis involves the mucosa of the large intestine, while Crohn's disease is marked by transmural inflammation and may affect any part of the gastrointestinal tract, but the majority of cases include the ileo-colon. IBD is a serious health issue because of its negative impact on a patient's quality of life and its high frequency, which has risen in recent years. Despite extensive research, the cause of IBD remains largely unexplained. Colitis is similar to the intestinal infection seen in IBD. *Capra hircus* milk oligosaccharides have been found to have anti-inflammatory properties.




Figure 4: Health benefits of Goat Milk.

The oligosaccharides avoid the anticipated decrease in body weight, increased colon size, and necrotic lesion extension (Figure 4). They also triggered a weakened immune response with less neutrophil penetration, as well as reduced clinical symptoms (diarrhea and bloody stools).^{16,36}

Malabsorption Disorders

Abnormalities in the absorption of food nutrients through the gastrointestinal tract cause malabsorption disorders. It may be affected by the side effects or complications of a variety of cancers. Malabsorption leads to deficiencies of a variety of nutrients, including Vitamin B₁₂, folic acid, and iron anemia, as well as other minerals, vitamins, and macronutrients. Malabsorption disorder is frequently concentrated in rodent models where the condition is incited by the response of about half of their small digestive tract. When looking at diets of *Capra hircus* and cow milk individually, examines showed improved stomach-related use of fat and protein, and higher evident edibility coefficient and retention of calcium, phosphorus, magnesium, iron, copper, zinc, and selenium. The higher protein content, cysteine levels, and amount of vitamin C and D in *Capra hircus* milk, relative to cow milk, is thought to be responsible for the better metabolic use of certain minerals.^{17,37}

Treatment of Cancer, Allergy and Others

Proteins are the most common antigens found in foods. Milk is one of the most common food allergens, and sensitivities to it are the most common in young children, with a 2-6 % frequency.¹⁸ Allergies can be acute or chronic, with symptoms ranging from non-life-threatening conditions like eczema, rhinitis, and intestinal issues to life-threatening conditions like anaphylaxis, bronchospasm, and urticaria. Due to its different protein composition, specifically its casein micelle elements, *Capra hircus* milk is a much less allergenic alternative to cow milk, according to several reports and anecdotal evidence.¹⁹ *Capra hircus* milk has shown critical enhancements in colic, minor stomach related issues, asthma, and skin inflammation over cow milk, just as in babies and kids with cow milk sharpens,²⁰ demonstrated that treatment with *Capra hircus* milk commonly settles somewhere in the range of 30 and 40% of issue instances of youth cow milk hypersensitivity, which can be higher sometimes. now and again outrage related unfavorably susceptible explanation, lactose prejudice is brought about by a lack in the lactase chemical used to process the milk sugar, lactose. Unhydrolyzed lactose travels through the large intestine of lactose-intolerant people, where it is fermented by bacteria, producing gases including hydrogen, methane,

carbon dioxide, and short-chain fatty acids, causing stomach problems including flatulence, abdominal discomfort, and diarrhea. Conjugated linoleic acid (CLA) is abundant in *Capra hircus* milk [74]. CLA has been shown to have anti-carcinogenic properties in animal models of mammary and colon cancer, as well as *in vitro* models of human melanoma,^{21,38} colorectal, and breast cancer. Although perturbation of eicosanoid-dependent cell signaling mechanisms, anti-oxidative effects, and disruption of estrogen receptor-mediated activities have all been suggested through fermented *Capra hircus* milk, the mechanism by which CLA inhibits tumor production is not fully known.²²

Prevention of Milk Allergy

Cow milk allergy is a prevalent condition in puberty and infancy, with an incidence of around 2.5 percent in the first three years of life. Cow milk is a vital nutrients source and is often one of the first forms of food proteins added to children. The infants are normally given cow milk-dependent formula after or in addition to breastfeeding. However, a replacement is needed for children who experience allergic symptoms shortly after ingestion. Cow milk allergy is an IgE-mediated allergy, which means that for unexplained causes, the body begins to develop IgE antibodies against specific proteins (allergens) in cow milk. When a human consumes milk regularly, an immune response occurs, resulting in allergic symptoms such as eczema, respiratory symptoms such as hay fever, wheezing, or asthma, gastrointestinal symptoms, or even anaphylaxis. In cow's milk allergy, the proteins α_{s1} -casein and β -lactoglobulin are essential allergens. Certain therapies, such as prolonged heating, can help to reduce the allergy-causing properties of β -lactoglobulin. Also, after a good denaturing operation, caseins retain the ability to bind to IgE.^{23,39} Cow milk proteins have a greater binding ability to IgE and IgG than *Capra hircus* milk proteins, according to an *in vitro* analysis. Cow milk has been shown to induce increased lymphocyte proliferation, IL-4 development, and histamine secretion and IgG production in animal models.²⁴

Alleviation of Lactose Intolerance

Lactose is used in *Capra hircus* milk, as well as cow and human milk. Despite this, *Capra hircus* milk is tolerated by many people with lactose intolerance. The superior digestibility of *Capra hircus* milk has been proposed as the cause. *Capra hircus* milk is ingested more thoroughly and quickly than cow milk, leaving less undigested waste in the colon to ferment and induce lactose intolerance symptoms. It's also possible that the patients don't have lactose sensitivity but are allergic to the main s1-casein proteins found in cow milk, which are usually lower or absent in *Capra hircus* milk. Lactose sensitivity and milk protein allergy have almost similar effects.^{12,40}

Anticarcinogenic Effect

Conjugated linoleic acid is abundant in *Capra hircus* milk. Conjugated linoleic acid has been shown to have anticarcinogenic properties in animal models of mammary and colon cancer, as well as *in vitro* models of human melanoma, colorectal, and breast cancer. While perturbation of eicosanoid-dependent cell signaling mechanisms, antioxidative effects, and disruption of estrogen receptor-mediated activities have all been proposed, the mechanism by which CLA inhibits tumor growth remains unknown.^{25,41}

Effect on Infancy Intake

In many parts of the world, feeding *Capra hircus* milk to infants has been and continues to be common. The scientific consensus on *Capra hircus* milk in child feeding is ambiguous. However, everyone agrees that unpasteurized *Capra hircus* milk cannot be supplied to babies and children because it is directly crucial to their survival. Unpasteurized

milk can contain pathogens or viruses that can cause life-threatening infections such as Q-fever, toxoplasmosis, tuberculosis, and brucellosis.²⁶ Pasteurized *Capra hircus* milk or formula based on *Capra hircus* milk may be a good substitute for cow milk. Commercially manufactured formulae are preferred since they are normally supplemented with folic acid, vitamin B12, and iron, which are deficient in daily *Capra hircus* milk. Infants who were fed homemade *Capra hircus* milk formulations and later diagnosed with megaloblastic anemia caused by folate and vitamin B12 deficiency have been reported in case studies.^{27,42}

Prevention of Inflammatory Bowel Disease (IBD)

Ulcerative colitis and Crohn's disease are two separate but related inflammatory bowel diseases. IBD is characterized by persistent and relapsing bowel inflammation, but it differs significantly from Crohn's disease in terms of pathophysiology and care. Ulcerative colitis includes the mucosa of the expansive digestive tract, whereas Crohn's illness is checked by transmural aggravation and may influence any portion of the gastrointestinal tract, but the larger part of cases incorporates the ileocolonic range. Despite extensive research, the cause of IBD remains largely uncertain.²⁸ *Capra hircus* milk's oligosaccharides have been found to have anti-inflammatory properties. The oligosaccharides avoid the anticipated decrease in body weight, increased colon size, and necrotic lesion extension. They also triggered a weakened immune response with less neutrophil penetration, as well as a reduction in clinical symptoms.^{29,43}

Immunomodulatory Activity

On the off chance that *Capra hircus* milk is not an ideal option for individuals with cow milk hypersensitivity, exceptionally late examinations showed invulnerable modulatory impacts from *Capra hircus* milk both *in vitro* and human studies, recently researched the impacts of *Capra hircus* milk on human platelets regarding nitric oxide (NO) and cytokine release.⁴⁴ The results exhibited that *Capra hircus* milk had the option to actuate NO delivery from platelets just as a set-off of cytokine creation (IL-10, TNF- and IL-6). The NO delivery could have cardio defensive impacts on the milk customer and uncover antibacterial action and consequently forestall diseases.³⁰ The most important participants in the innate and adaptive immune responses are T-lymphocytes (T-cells), Natural Killer (NK) cells, and B-lymphocytes (B-cells). Although the composition of immunoglobulins (Ig) is similar, small variations among the main immunological groups (IgG, IgM, IgA, IgD, and IgE) are linked to several biological properties, and IgG and IgA account for the majority of serum immunoglobulins. *Capra hircus* milk is involved in nearly all biochemical processes and has anti-inflammatory and antioxidant properties. This can be critical as aggravation is the body's essential reaction to contamination and oxidation has been connected to the advancement of numerous illnesses, counting cancer. Besides, other variables such as the maintenance of a solid intestinal microflora with the assistance of probiotics and prebiotics are fundamental for ensuring against the negative impacts of pathogenic contamination sensitivity.^{31,41}

CONCLUSION AND FUTURE PROSPECTS

Capra hircus milk and other *Capra hircus*-derived products have distinct properties, and their nutritional importance, as well as their possible health effects, have been studied extensively. The composition of *Capra hircus* milk is similar to that of cow milk, but it is more similar to human milk, making it more tolerable. Furthermore, *Capra hircus* milk's superior digestibility, fatty acid content, and inclusion of different bioactive compounds in its composition make it potentially beneficial in the treatment or even prevention of certain medical conditions. Anti-inflammatory and anti-oxidative properties of *Capra hircus* milk, whey,

and fermented *Capra hircus* products can reduce the risk of chronic diseases. *Capra hircus* milk and its subordinates may too advance the particular improvement of microscopic organisms that are a portion of the intestinal microbiota, which can have metabolic, endocrine, and resistant framework benefits.⁴⁵ In combination with traditional medical therapy, *Goat's* milk and other *Capra hircus* derivatives can serve as health-promoting nutrients and increase overall clinical performance in the management of chronic diseases. Infants, the sick, and those recovering from illness should drink *Capra hircus* milk. Studies are needed to reduce *Capra hircus* milk's "Goaty taste," which makes it less desirable.⁴³ According to animal studies, *Capra hircus* milk can help with malabsorption and inflammatory bowel disease. Fermented *Capra hircus* milk has anti-oxidative anti-atherogenic properties, which can diminish the chance of cardiovascular illness.⁴⁸ However, further human trials are required before *Capra hircus* milk products can be said to have health benefits.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

REFERENCES

- Mahmoud SA, Daud Ali M, Ahmad A, Maghrabi AA, Harthi MA, Fattani NA. Knowledge and practice about use of medication among breast feeding women in Saudi Arabia: A prospective cohort study. *Ijpi*. 2019;9(1):16-9. doi: 10.5530/ijpi.2019.1.5.
- Roncada P, Gaviraghi A, Liberatori S, Canas B, Bini L, Greppi GF. Identification of caseins in goat milk. *Proteomics*. 2002;2(6):723-6. doi: 10.1002/1615-9861(200206)2:6<723::AID-PROT723>3.0.CO;2-I, PMID 12112854.
- Goswami M, Bharti SK, Tewari A, Sharma H, Karunakara KN, Khanam T. Implication of functional ingredients of goat milk to develop functional foods. *Anim Feed Sci Technol*. 2017;5:65-72.
- Manso MA, Escudero C, Alijo M, López-Fandiño R. Platelet aggregation inhibitory activity of bovine, ovine, and caprine κ -casein macropolypeptides and their tryptic hydrolysates. *J Food Prot*. 2002;65(12):1992-6. doi: 10.4315/0362-028x-65.12.1992, PMID 12495023.
- Qian ZY, Jollès P, Migliore-Samour D, Schoentgen F, Fiat AM. Sheep κ -casein peptides inhibit platelet aggregation. *Biochim Biophys Acta (BBA) Gen Subj*. 1995;9:411-7.
- Boyazoglu J, Morand-Fehr P. Mediterranean dairy sheep and goat products and their quality. A critical review. *Small Rumin Res*. 2001;40(1):1-. doi: 10.1016/S0921-4488(00)00203-0, PMID 11259871.
- Bevilacqua C, Martin P, Candali C, Fauquant J, Piot M, Roucayrol AM *et al*. Goats' milk of defective α 1-casein genotype decreases intestinal and systemic sensitization to β -lactoglobulin in guinea pigs. *J Dairy Res*. 2001;68(2):217-27. doi: 10.1017/S0022029901004861.
- Assis POAd, Guerra GCB, Araújo DFdS, Araújo Júnior Rfd, Machado TADG, Araújo AAd *et al*. Intestinal anti-inflammatory activity of goat milk and goat yoghurt in the acetic acid model of rat colitis. *Int Dairy J*. 2016;56:45-54. doi: 10.1016/j.idairyj.2015.11.002.
- López-Aliaga I, Díaz-Castro J, Alférez MJ, Barrionuevo M, Campos MS. A review of the nutritional and health aspects of goat milk in cases of intestinal resection. *Dairy Sci Technol*. 2010;0(6):611-22.
- Park YW. Hypo-allergenic and therapeutic significance of goat milk. *Small Rumin Res*. 1994;14(2):151-9. doi: 10.1016/0921-4488(94)90105-8.
- Lima MJ, Teixeira-Lemos E, Oliveira J, Teixeira-Lemos LP, Monteiro A, Costa JM. Nutritional and health profile of goat products: focus on health benefits of goat milk. *Goat Sci*. *Intech Open*. 2018;20:189-232.
- Park YW, Juárez M, Ramos M, Haenlein GFV. Physico-chemical characteristics of goat and sheep milk. *Small Rumin Res*. 2007;68(1-2):88-113. doi: 10.1016/j.smallrumres.2006.09.013.
- Jenness R. Composition and characteristics of goat milk: Review 1968–1979 [review]. *J Dairy Sci*. 1980;63(10):1605-30. doi: 10.3168/jds.S0022-0302(80)83125-0.
- Yadav AK, Singh J, Yadav SK. Composition, nutritional and therapeutic values of goat milk: a review. *Asian J Dairy Food Res*. 2016;35(2):96-102. doi: 10.18805/ajdr.v35i2.10719.
- Zenebe T, Ahmed N, Kabeta T, Kebede G. Review on medicinal and nutritional values of goat milk. *Acad J Nutr*. 2014;3(3):30-9.
- Cerbulis J, Parks OW, Farrell Jr HM. Composition and distribution of lipids of goats' milk. *J Dairy Sci*. 1982;65(12):2301-7. doi: 10.3168/jds.S0022-0302(82)82501-0.
- Jensen RG, Ferris AM, Lammi-Keefe CJ, Henderson RA. Lipids of bovine and human milks: a comparison. *J Dairy Sci*. 1990;73(2):223-40. doi: 10.3168/jds.S0022-0302(90)78666-3, PMID 2184172.
- Ceballos LS, Morales ER, de la Torre Adarve G, Castro JD, Martínez LP, Sampelayo MRS. Composition of goat and cow milk produced under similar conditions and analyzed by identical methodology. *J Food Compos Anal*. 2009;22(4):322-9. doi: 10.1016/j.jfca.2008.10.020.
- Slacanac V, Božanic R, Hardi J, Rezessyné Szabó JU, Lucan M, Krstanovic V. Nutritional and therapeutic value of fermented caprine milk. *Int J Dairy Technol*. 2010;63(2):171-89. doi: 10.1111/j.1471-0307.2010.00575.x.
- Tomotake H, Okuyama R, Katagiri M, Fuzita M, Yamato M, Ota F. Comparison between Holstein cow's milk and Japanese-Saanan goat's milk in fatty acid composition, lipid digestibility and protein profile. *Biosci Biotechnol Biochem*. 2006;70(11):2771-4. doi: 10.1271/bbb.60267, PMID 17090948.
- Díaz-Castro J, Hijano S, Alférez MJ, López-Aliaga I, Nestares T, López-Frías M *et al*. Goat milk consumption protects DNA against damage induced by chronic iron overload in anaemic rats. *Int Dairy J*. 2010;20(7):495-9. doi: 10.1016/j.idairyj.2010.01.006.
- Almaas H, Cases AL, Devold TG, Holm H, Langsrud T, Aabakken L *et al*. *In vitro* digestion of bovine and caprine milk by human gastric and duodenal enzymes. *Int Dairy J*. 2006;16(9):961-8. doi: 10.1016/j.idairyj.2005.10.029.
- Atanasova J, Ivanova I. Antibacterial peptides from goat and sheep milk proteins. *Biotechnol Biotechnol Equip*. 2010;24(2):1799-803. doi: 10.2478/V10133-010-0049-8.
- Horácková Š, Sedláčková P, Sluková M, Plocková M. The influence of whey, whey component and malt on the growth and acids production of lactobacilli in milk. *Czech J Food Sci*. 2014;32(6):526-31. doi: 10.17221/214/2014-CJFS.
- Kunz C, Rudloff S, Baier W, Klein N, Strobel S. Oligosaccharides in human milk: structural, functional, and metabolic aspects. *Annu Rev Nutr*. 2000;20(1):699-722. doi: 10.1146/annurev.nutr.20.1.699, PMID 10940350.
- Lara-Villoslada F, Debras E, Nieto A, Concha A, Gálvez J, López-Huertas E *et al*. Oligosaccharides isolated from goat milk reduce intestinal inflammation in a rat model of dextran sodium sulfate-induced colitis. *Clin Nutr*. 2006;25(3):477-88. doi: 10.1016/j.clnu.2005.11.004, PMID 16375993.
- Daddaoua A, Puerta V, Requena P, Martínez-Férez A, Guadix E, de Medina FS *et al*. Goat milk oligosaccharides are anti-inflammatory in rats with hapten-induced colitis. *J Nutr*. 2006;136(3):672-6. doi: 10.1093/jn/136.3.672, PMID 16484541.
- Hernández-Ledesma B, Ramos M, Gómez-Ruiz JÁ. Bioactive components of ovine and caprine cheese whey. *Small Rumin Res*. 2011;101(1-3):196-204. doi: 10.1016/j.smallrumres.2011.09.040.
- Bhattarai RR. Importance of goat milk. *J Food Sci Technol Nepal*. 2014;7:107-11. doi: 10.3126/jfstn.v7i0.11209.
- Csapóné Riskó T, Csapó Z. Goat keeping and goat milk products in human nutrition - review. *APSTRACT*. 2019;13(1-2):24-36. doi: 10.19041/APSTRACT/2019/1-2/3.
- Turkmen N. Chapter 35 – the nutritional value and health benefits of goat milk components, nutrients in dairy and their implications on health and disease. *Academic Press*; 2017. p. 441-9.
- Sepe L, Argüello A. Recent advances in dairy goat products. *Asian-Australas J Anim Sci*. 2019;32(8):1306-20. doi: 10.5713/ajas.19.0487, PMID 31357271.
- Lad SS, Aparnathi KD, Mehta B, Velpula S. Goat milk in human nutrition and health – a review. *IntJCurrMicrobiolAppSci*. 2017;6(5):1781-92. doi: 10.20546/ijcmas.2017.605.194.
- Kumar S, Kumar B, Kumar R, Kumar S, Khatkar SK, Kanawjia SK. Nutritional F nutritional features of goat milk es of goat milk- A review. *Indian J Dairy Sci*. 2012;65(4):266-73.
- Roy D, Ye A, Moughan PJ, Singh H. Composition, structure, and digestive dynamics of milk from different species- A review. *Front Nutr*. 2020;7:577759. doi: 10.3389/fnut.2020.577759, PMID 33123547.
- Vaquil, Rekha R. A review on health promoting aspects of goat milk. *The Pharm Inno*;J2017;6(12):05-8.
- Filipczak-Fiutak M, Pluta-Kubica A, Domagala J, Duda I, Migdał W. Nutritional value and organoleptic assessment of traditionally smoked cheeses made from goat, sheep and cow's milk. *PLOS ONE*. 2021;16(7):e0254431. doi: 10.1371/journal.pone.0254431, PMID 34293016.
- Lin X, Luo J, Zhang L, Wang W, Gou D. MiR-103 controls milk fat accumulation in goat (*Capra hircus*) mammary gland during lactation. *PLOS ONE*. 2013;8(11):e79258. doi: 10.1371/journal.pone.0079258, PMID 24244462.
- Marrone R, Ramkumar A, Smaldone G, Ruffano D, Chirollo C, Veneziano V *et al*. Deltamethrin residues in milk and cheese of lactating goats (*Capra hircus*). *Molecules*. 2019;24(3):517. doi: 10.3390/molecules24030517, PMID 30709021.
- Moura RR, Melo LM, Freitas VJdF. Production of recombinant proteins in milk of transgenic and non-transgenic goats. *Braz Arch Biol Technol*. 2011;54(5):927-38. doi: 10.1590/S1516-89132011000500010.
- Collins RA, Harper AE, Schreiber M, Elvehjem CA. The folic acid and vitamin B₁₂ content of the milk of various species. *J Nutr*. 1951;43(2):313-21. doi: 10.1093/jn/43.2.313, PMID 14851047.
- Karekar P, Killedar S. Efficient *Euphorbia hirta* Phytosomes for *in vitro* antiasthmatic activity. *IJPI*. 2022;12(2):178-82. doi: 10.5530/ijpi.2022.2.32.

43. Van Leeuwen SS, Te Poele EM, Chatziioannou AC, Benjamins E, Haandrikman A, Dijkhuizen L. Goat Milk Oligosaccharides: Their Diversity, Quantity, and Functional Properties in Comparison to Human Milk Oligosaccharides. *J Agric Food Chem.* 2020;68(47):13469-85. doi: 10.1021/acs.jafc.0c03766, PMID 33141570.
44. Kapadiya DB, Prajapati DB, Jain AK, Mehta BM, Darji VB, Aparnathi KD. Comparison of Surti goat milk with cow and buffalo milk for gross composition, nitrogen distribution, and selected minerals content. *Vet World.* 2016;9(7):710-6. doi: 10.14202/vetworld.2016.710-716, PMID 27536031.
45. Zobel G, Neave HW, Webster J. Understanding natural behavior to improve dairy goat (*Capra hircus*) management systems. *Transl Anim Sci.* 2019;3(1):212-24. doi: 10.1093/tas/txy145, PMID 32704793.
46. Karami S, Roayaei M, Hamzavi H, Bahmani M, Hassanzad-Azar H, Leila M *et al.* Isolation and identification of probiotic *Lactobacillus* from local dairy and evaluating their antagonistic effect on pathogens. *Ijpi [internet].* Vol. 7(3). p. 137-41; 2020 [cited 12/9/2022].

Article History: Submission Date : 21-06-2022; Revised Date : 04-07-2022; Acceptance Date : 15-08-2022.

Cite this article: Tiwari G, Chauhan A, Sharma P, Tiwari R. Nutritional Values and Therapeutic uses of *Capra hircus* Milk. *Int. J. Pharm. Investigation.* 2022;12(4):408-17.